References:

Ahluwalia S.S.;Goya D. (2007), Microbial and plant derived biomass for removal of heavy metals and plants, *FEMS Microbial Rev*, 25; 335–347.

Asgari A. R.; Vazi F.; Nasseri S.; Dordelmann O.;Mahvi A. H.; Fard E. D. (2008), Removal of hexavalent chromium from drinking water by granular ferric hydroxide, *Iran. J. Environ. Health.SCi. Eng*,5(4); 277-282.

ATSDR (Agency for Toxic Substances and Disease Registry), (1989), Toxicological profilefor chromium, Prepared by Syracuse Research Corporation under Contract 68-C8-0004.Syracaue, USA U.S.,*Public Health Service, ATSDR/TP*-88/10.

Baral A.; Engelken.R.; Stephens.W.; Farris.J.; Hannigan.R. (2006). Evaluation of aquatic toxicities of Chromium and chromium- containing effluent in reference to chromium electroplating industries, *Arch.Environ.Contam. Toxicol*, 50; 496-502.

Baral A. and Engelken. R. (2002), Chromium – based regulation and greening in metal finishing industries in the USA, *Environment Sci. Policy*, 52; 121-133.

Bartlett R.J. and James B.R (1988), Mobility and bioavailability of chromium in soils.*In `Chromium in the natural and human environments, (Eds JO Niragu, E Nieboer),(JohnWiley and Sons Inc.: New York);* 267-303.

Benhammou A.; Yaacoubi A.; Nibou L.; Tonouti B. (2005), Study of the removal of Hg(II) and Cr(VI) from aqueous solution by Moroccan stevensite, *J. Hazardous Materia*, 117; 243-249.

Beszedits. Sci.(1998), Chromium removal from industrial waste waters, in: J.O.Nriagu, E. Nieboer (Eds), *Chromium in the Natural and Human Environments, John Wiley, New York*; 232-263.

Bruins, M. R.; Kapil, S.; Oehme F. W. (2000), Microbial resistance to metals in the environment, *Ecotox. Environ. Safe*, 45; 198-207.

Cervantes.C.; Campos-Garcıa J.;Devars.S.; Gutierrez-Corona F.;Loza-Tavera H.; Torres-Cervantes C.; Campos-Garcia.J. ;Devars. S.; Gutierrez-Corona F.; Loza-TaveraH.;Torres-Guzman J.C.; Moreno-Sanhez.R.(2001), Interaction of chromium with microorganisms and plants, *FEMS Microbial Rev*, 25; 335-347.

Das A. and Mishra S. (2008),Hexavalent Chromium (VI): Health hazards &Environmental pollution,*Journal of Environment Research and Development*, 2; 386-392.

Donmez G.; Aksu Z. (2002), Removal of Cr(VI) from saline waste water by Dunaliellaspecies, *Proc. Biochem*, 38; 751-762.

FloresA.and Perez J. M. (1999), Cytotoxicity, apoptosis, and in vitro DNA damage induced by potassium chromate, toxicol, Appl. Pharmocol,*Prepared by Syracuse Research Corporation under Contract 68-C8-0004. Syracaue, USA U.S*, 161; 75-81.

Gauglhofer J. and Bianchi V. (1991), Chromium. In: Merian E (Ed), Metals and their compounds in the environment,*New York: VCH Publisher*; 853-878.

Guzman. J. C.; Moreno-Sanchez R. (2001), Interactions of chromium with microorganismsLarsJärup Hazards of heavy metal contamination,British Medical Bulletin;**68**; 167–182.

Hesham M. A.; Engy M. K.;Amr H. M.; Ahmed D.(2010), Chromium removal from annery waste water using chemical and biochemical techniques aming zero discharge of pollution, *Proceeding of fifth Scientific Environmental Conferance, Zagazig University*; 171-183.

Kar D.; Sur P.;Mandal S. K.; Saha T.; Kole R. K. (2008), Assessment of heavy metal pollution in surface waterInt, *J. Environ. Sci. Tech*., 5 (1); 119-124.

Kotas.J. and Stasicka. Z. (2000), Chromium occurrence in the environment and methods of its speciation,*Environment pollution*, 107; 263-283.

Lofroth. G. and Ames B.N. (1978), Mutagenicity of inorganic compounds in Salmonella typhimurium: arsenic, chromium and selenium,*Mutat.Res*, 53; 65-66.

Mahmood M. B.;Nagham A. A.; Manal A. T. (2008), Removal Of Chromium From Electro-Plating Wastewater By Simple Chemical Treatment And Ion Exchange, *Eng and Tech*, 26(11); 212-215.

Malik A, (2004), Metal bioremediation through growing cells,*Environment International*, 30; 261-278.

Marschner H. (1995), Mineral nutrition of higher plants,Academic Press, London.Matzat E. and Shiraki K (1974), Chromium, *Handbook of Geochemistry*, 11(3); 24(A).

McGrath S.P. and Smith.S. (1990), Chromium and Nickel, in B.J.Alloway (ed), *Heavy Metals in soils, John Wiley and Sons, New York*; 125-147.

Middleton. S.S.; Latmani R. B.; Mackey M. R.; Ellisman M. H.; Tebo. B.M.; Criddle C.S. (2003), Cometabolism of Cr (VI) by *Shewanellaoneidensis*MR-1 produces cell-associated reduced chromium and inhibits growth, *Biotechnol. Bioeng*, 83; 627–637.

Mirlean N.; Rosenberg A.; Chies JO (2007), Metal contamination of vineyard soils in wet subtropics (southernBrazil), *Environmental Pollution***,**149; 10-17.

Mohan D.; Pittman U. C. Jr. (2006), Activated carbons and low cost adsorbents forremediation of tri- and hexavalent chromium from water, *Journal of hazardous Materials*, B137; 762-811.

Mohhapatra P. K. (2006), Textbook of environmental biotechnology, *I. K. International publishing house, Mumbai*, 43-51.

Moncur M.C.;Ptacek C.J.; Blowers D.W.; Jambor J. L. (2005), Release, Transport, and attenuation of metal from an old tailing impoundment, *Applied Geochemistry*, 20; 639-659.

Montanher S. F.; Oliveria E. A.; Rollemberg M. C.; (2005), Removal of metal ions from aqueous solutions by sorption on to rice barn, *J. Hazardous material*, B 1117; 207-211.

Moore J.W. and Ramamurthy S. (1984), Heavy Metg. Neals in Nature Waters,*Springer-Verla. New York*, 28(57); 182-184.

Namasivayam C. and Ranganathan K. (1993),Waste Fe (III)/Cr(III) hydroxide as adsorbent for the removal of Cr(VI) from aqueous solution and chromium plating industry wastewate, *Environmental Pollution*, 82; 255-261.

Ngah W. W.; Endud C. S.;Mayanar R. (2002), Removal of copper (II) fromaqueous solution onto chitosan and cross-linked chitosan beads, *React FunctPolym*, 50; 181-90.

Nishioka H. (2009), Mutagenic activity of metal compounds in bacteria,*Mutation Research*, 31; 185-189.

Nriagu J. O.; Pacyna J.M. (1988), Quantitative assessment of worldwide contamination of air, water, and soils by trace metals,*Nature*, 333; 134-139.

Pandi M.; Shashirekha V.; Swamy M. (2007), Biosorptionof chromium from retain chrome liquor by cyano bacteria, Microbiological research, 1-9metals from wastewater, *BioresourTechnol*, 98; 2243–2257.

Richard F.C. and Bourg C.M (1991), Aqueous Geochemistry of chromium: A Review, *Water Research*, 10; 824-827.

Robles-Camacho J.;Armienta M.A. (2000), Natural chromium contamination of groundwater at Leon valley, *Mexico. Journal of Geochemical Exploration*, 68; 167-181.

Ryan M. P.; Williams D.E.; Chater R. J.; Hutton B.M.; McPhail D. S. (2002), Why stainlss steel corrodes?, *Nature*, 415; 770-774.

Ryan M.P.; Williams D. E.; Chate R. J.; Hutton B. M.; McPhail D. S. (2002),Why stainlesssteel corrodes? , *Nature*, 415; 770-774.

Salnikow K. and Zhitkovich. A. (2008), Genetic and Epigenetic Mechanisms in MetalCarcinogenesis and Co carcinogenesis: Nickel, Arsenic, and Chromium, *Chem. Res. Toxicol*, 21; 28-44.

Sedlak D.L. and Chan P.G. (1997), Reduction of hexavalent chromium by ferrous iron,*Geochemical et.al.CosmochimicaActa*, 61; 2185-2192.

Sharma D.C.; ChatterjeeC.; Sharma C. P. (1995), Chromium accumulation and its effects on wheat (Triticumaestivum L., cv. HD 2204) metabolism,*Plant Science*, 111; 145-151.

SikailyEi.;Nemr A. EI; Khaled A. (2007), Removal of toxic chromium from waste water,*Indian Journal of Technology*, 15; 19-28.

Sujan M.; Gude;Das S. N. (2008), Adsorption of Cr(VI from aqueous solution by chemically treated water by hyacinth Eichhorniacrassipes,*Indian Journal of Technology*, 15; 12-18.

United State Environmental Protection Agency (USEPR), (1998), Toxicological review of hexavalent chromium (CAS No.18540-29-9), USEPA, Washington DC.

United States Environmental Protection Agency (US EPA), (1998), Toxicological review ofusing green alga Ulvalactuca and its activated carbon, *J. Hazard. Mater*, 148 2; 16-228.

Venitt S. L. (1994), Mutagenicity of chromate in bacteria and its relevance to chromate carcinogenesis, *Nature*, 250; 493-495.

Viamajala S. P.;SaniB.M.; ApelR.K.,; Petersen W.A. (2004), Toxic effect of Chromium (VI) on Anaerobic and Aerobic Growth of Shewanellaoneidensis,*MR-1. Biotechnol, Prog*, 20; 87-95.